

## **NEW**

# PROBIOTIC SUPPLEMENTARY **FEED FOR POULTRY**

## CHICK'A

- Probiotic supplementary feed for poultry
- Improves poultry welfare
- Improves egg-laying
- Stabilises intestinal flora
- Suitable for all poultry (chickens, pullets, breeding hens, turkeys, etc.)

Composition: Corn Starch, Brewer's Yeast, Inulin, Calcium Carbonate

### Ingredients:

- Raw protein: 10,2 %
- Raw fat <1.0 %</li>
- Fibers < 0.3 %
- Ash 22,8 %
- Humidity 6,0 %

**Dosage**: 360 g, equivalent to 48 days' care for 5 hens







**UKAL.**com









#### **ZOOTECHNICAL ADDITIVES:**

intestinal flora stabilizer Bacillus velezensis (4b1820) 3 x 1011 CFU/kg

### NUTRITIONAL PHYSIOLOGICAL ADDITIVES:

Calcium 80.3 g/kg, Magnesium 600 mg/kg, Potassium 4500 mg/kg, Sodium 200 mg/kg, Phosphorus 2400 mg/kg, Vitamin D (3a670a) 500 IU/kg, vitamin E (3a700) 20 mg/kg, vitamin K3 (3a711) 0.62 mg/kg, vitamin B1 (3a821) 0.5 mg/kg, vitamin B2 (3a825ii) 1.25 mg/kg, vitamin PP (3a315) 8.75 mg/kg, pantothenic acid (3a841) 3.75 mg/kg, vitamin B6 (3a831) 1.25 mg/kg, biotin (3a880) 25 mcg/kg, vitamin B9 (3a316) 0.4 mg/kg, vitamin C (3a300) 25 mg/kg, iodine (3b202) 0.5 mg/kg, copper (3b410) 5 mg/kg, manganese (3b510) 15 mg/kg, zinc (3b610) 10 mg/kg, iron 130 mg/kg

#### **FEEDING ADVICE:**

Give 12g / kg feed over the feed per day. Equivalent to 1.5g per day for a 2.5kg chicken with 120g feed. The enclosed measuring spoon holds 7.5g, which corresponds to 5 servings (5 days / chicken or 1 daily dose for 5 chickens). If necessary, increase dosage by 1 measuring spoon.

**Suitable for permanent use.** An optimal result is achieved with regular and long-term use.

#### **SHELF LIFE:**

2 years from production date. Store dry at room temperature.

